arXiv:1207.0646v1 [cond-mat.str-el] 3 Jul 2012

Anomalous Hall effect in heavy electron materials

Yi-feng Yang
Beijing National Laboratory for Condensed Matter Physics and
Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China
(Dated: July 4, 2012)

We propose a new empirical formula for the anomalous Hall effect in heavy electron materials based on a phenomenological two-fluid description of the f-electron states. The new formula incorporates two previous theories proposed by Fert and Levy in 1987 and Kontani and Yamada in 1994 and takes into account both incoherent and coherent skew scatterings from local and itinerant f-electrons. We perform experimental analysis in several heavy electron compounds and show that the new formula provides a consistent description of the evolution of the Hall coefficient in the whole temperature range down to only a few Kelvin.

PACS numbers: 71.27.+a, 72.10.-d, 72.15.Gd

Anomalous Hall effect has attracted much interest in recent years due to its topological origin [1]. In general, the measured Hall coefficient R_H includes two terms, $R_H = R_0 + R_s$, where R_0 is the ordinary Hall coefficient and R_s is the extra-ordinary or anomalous Hall coefficient. The microscopic origin of R_s has proved quite complicated. Three distinct contributions – intrinsic [2], skew scattering [3] and side-jump [4] – have been identified; each of them has an individual scaling, $R_s \propto \rho^\alpha$, with respect to the longitudinal resistivity ρ. In ferromagnetic conductors, a simple summation of the three terms yields an empirical formula that explains a large amount of experimental data.

In heavy electron materials, however, such an empirical formula is not available despite a number of theoretic proposals [5–9]. So far the most prevailing theory was developed by Fert and Levy in 1987 [6]. They considered incoherent skew scattering of conduction electrons by independent f-moments and predicted,

$$R_s = r_l \rho \chi, \tag{1}$$

where χ is the magnetic susceptibility and r_l is a constant. Their theory has been verified in $CeAl_3$, $CeCu_2Si_2$ and most other materials in the high temperature regime but fails when coherence sets in. In the caged compound $Ce_3Rh_4Sn_{13}$, in which no lattice coherence is observed, the scaling persists down to the lowest measured temperature [10]. In most nonmagnetic Ce- and U-compounds such as $CeRu_2Si_2$, CeNi, $CeCu_6$, UPt_3 and UAl_2, a different scaling, $R_s \propto \rho^2$, has been observed at very low temperatures and explained by the coherent skew scattering of f-electrons [7–9]. In the nonmagnetic compound Ce_2CoIn_8, both formulas seem to apply in their respective high or low temperature regime [11].

But unlike in ferromagnetic conductors, a direct summation or interpolation of the two contributions cannot explain the experimental result in the intermediate temperature regime. The theory of Kontani and Yamada [9] extrapolates to a quite different scaling,

$$R_s = r_h \chi, \tag{2}$$

where r_h is a constant, and necessarily fails to describe localized f-moments at high temperatures. The fundamental difference in dealing with f-electrons makes the combination of the two theories, if possible at all, highly nontrivial, which requires detailed knowledge about the incoherent and coherent behaviors of the f-electrons. Thanks to the recent development of a phenomenological two-fluid framework [12], we now have a quantitative measure of the temperature evolution of the f-electron states. In this work, we apply the two-fluid description to the intermediate temperature regime and propose a new empirical formula for the anomalous Hall effect in heavy electron materials. We then perform data analysis in several compounds and present experimental evidences that provide unambiguous support to our formula.

Anomalous Hall effect in Ce-115 ($CeMIn_5$, M = Co, Ir, Rh) provides the first insight into this complicated problem. Fig. 1(a) reproduces the measured Hall coefficient in all three Ce-115 materials [13, 14]. The con-



FIG. 1: (color online) Hall coefficient of Ce-115 and Ce-218 materials reproduced from Refs. [13] and [15]. The incoherent skew scattering contribution is completely suppressed and the temperature dependence of the Hall coefficient does not follow any combination of $\rho^\alpha\chi$. Instead, it scales with the susceptibility of the itinerant Kondo liquid (solid lines) [16].

stant behavior at high temperatures indicates that the incoherent skew scattering contribution predicted by Fert and Levy is negligible small, namely $r_l \approx 0$. We can thus disregard the effect of localized f-moments and focus on the contribution of coherent f-electrons. It turns out that the Hall coefficient R_H cannot fit to any combination of $\rho^\alpha \chi$. Instead, within the framework of a phenomenological two-fluid scenario, it is found to scale with a universal fraction of the magnetic susceptibility, namely the susceptibility of an itinerant heavy electron Kondo liquid [16], $\chi_h \propto f_h(T)[1 + \ln(T^*/T)]$, where $f_h(T) = f_0(1 - T/T^*)^{3/2}$ is the fraction of the f-electron spectral weight that participates in the Kondo liquid formation. f_0 represents a hybridization effectiveness that controls the eventual itinerancy of the f-electrons [17].

The universal χ_h-scaling in the anomalous Hall effect of itinerant f-electrons is similar to the prediction of Kontani and Yamada at high temperatures [9], but only subject to its own magnetization given by χ_h. It is therefore expected that the two components contribute independently to the Hall effect, with conduction electrons deflected in each channel by a local field produced by the corresponding component. Taking together the incoherent contribution from localized f-moments [6], we arrive at

$$R_H = R_0 + R_s^l + R_s^h \approx R_0 + r_l \rho \chi_l + r_h \chi_h, \qquad (3)$$

where χ_l is the magnetic susceptibility of the residual unhybridized f-moments and $R_s^{l,h}$ are the anomalous Hall coefficient contributed by localized and itinerant f-electrons, respectively. We have made the simplification $R_s^h \approx r_h \chi_h$ for the coherent contribution since we are most interested in the intermediate temperature regime. Its crossover from ρ^0-scaling to ρ^2-scaling is not universal. Regardless of the details, we have on quite general basis, $R_s^h \sim \chi_h \gamma_h^2/(E_d^2 + \gamma_h^2)$, where the resonance energy E_d is typically of the order of T^*, while the broadening or hybridization γ_h increases from a few meV at zero temperature to $\sim 100\,\mathrm{meV}$ near T^* [18–20]. This gives $R_s^h \propto \rho^0$ for $\gamma_h \gg E_d$ at high temperatures and $R_s^h \sim \gamma_h^2 \propto \rho^2$ at very low temperatures if $\gamma_h < |E_d|$. The quite unusual form of R_H in Eq. (3) reflects the fundamental difference of heavy electron materials from ferromagnetic conductors in that, with lowering temperature, localized f-moments gradually dissolve into the Kondo lattice and may not be treated as a static spin polarized background. This is in fact a general feature of strongly correlated d-electron and f-electron systems and should always be taken into consideration [21].

Eq. (3) provides a new empirical formula for analyzing the Hall effect in heavy electron materials. As will be confirmed below, it tracks the whole temperature evolution of the Hall coefficient down to only a few Kelvin. The key of our analysis is to separate the magnetic susceptibility χ_l and χ_h so that the new formula can be applied to the experimental data in a straightforward manner. The scaling form $R_H = R_0 + r_h \chi_h$ in the limit $R_s^l \approx 0$ in all three Ce-115 compounds under pressure and in La-doped CeCoIn$_5$ has already been discussed in detail in [16] and will not be repeated here.



FIG. 2: (color online) Analysis of the Hall coefficient of URu$_2$Si$_2$ [18]. The ordinary Hall coefficient is $R_0 = -7.27 \times 10^{-3}\,\mathrm{cm}^3\mathrm{C}^{-1}$. We have $T^* = 65\,\mathrm{K}$ and $T_L = 10\,\mathrm{K}$. Using the new formula, we find a good fit in the whole temperature range above T_{HO}. The inset plots the two components of the susceptibility derived from the Knight shift analysis [17, 22, 23]. The derived χ_h follows exactly the Kondo liquid scaling (solid line).

We take URu$_2$Si$_2$ as the first example, in which Knight shift experiment [22, 23] allows us to determine χ_l and χ_h unambiguously. The two-fluid descriptions of the magnetic susceptibility and the Knight shift [16, 24, 25] read, $\chi = \chi_l + \chi_h$ and $K = K_0 + A\chi_l + B\chi_h$, where A and B are the hyperfine couplings. At high temperatures, f-electrons are well localized so that $K = K_0 + A\chi$ for $T > T^* \approx 65\,\mathrm{K}$; while at very low temperatures deep within the hidden order phase [26], all $5f$-electrons become itinerant and we have $K = K_0 + B\chi$ for $T < T_L \approx 10\,\mathrm{K}$. These determine the values of K_0, A and B without arbitrariness. Following the discussion in [17, 23], the two components χ_l and χ_h for $T_L < T < T^*$ are then immediately derived by using $\chi_l = (K - K_0 - B\chi)/(A - B)$ and $\chi_h = (K - K_0 - A\chi)/(B - A)$. The results are plotted in the inset of Fig. 2. As we can see, the partial susceptibility χ_h (diamond) falls exactly upon the scaling formula (solid line) of the Kondo liquid predicted in the two-fluid framework [16].

We now turn back to the Hall experiment in URu$_2$Si$_2$ [18]. In the high temperature regime $T > T^*$, all f-electrons are localized and, by using the experimental data for the magnetic resistivity [18], the Hall coefficient is found to scale very well with the prediction of Fert and Levy up to the highest measured temperature of about 300 K,

$$R_H = R_0 + r_l \rho \chi, \qquad (4)$$

which determines the value of R_0 and r_l. We then apply



FIG. 3: (color online) Comparison between the subtracted coherent contribution $R_H - R_0 - r_l\rho\chi_l$ and $\rho^\alpha\chi_h$ for $\alpha = 0$, 1, 2 in URu$_2$Si$_2$. Only for $\alpha = 0$, an overall proportionality is found in the whole temperature range between T^* and T_{HO}.

Eq. (3) to $T < T^*$ and note that there is only one adjusting parameter, r_h. The result is presented in Fig. 2 and a good fit is obtained in the whole temperature range above T_{HO}. This undoubtedly confirms our proposed new formula, especially considering that all parameters are determined uniquely by experiment without any adjustment. In order to verify other possible forms of the coherent contribution R_s^h, Fig. 3 compares $R_H - R_0 - r_l\rho\chi_l$ and $\rho^\alpha\chi_h$ calculated from experimental data for different values of α. Only for $\alpha = 0$ we find an overall agreement for temperatures between T_{HO} and T^*. For $T < T_{HO}$, different values of R_0 and r_h are required to fit the data, indicating a Fermi surface change across the hidden order transition [26].

Our analysis can be applied straightforwardly to other materials as long as χ_l and χ_h can be separated in experiment. Otherwise, a less rigorous method may be applied by using the scaling formula a prior [16],

$$\chi_h = \min\left\{\chi, \chi_0\left(1-\frac{T}{T^*}\right)^{3/2}\left(1+\ln\frac{T^*}{T}\right)\right\}, \qquad (5)$$

where χ is the experimental susceptibility and χ_0 is a constant prefactor related to the Wilson ratio of the heavy electrons and the hybridization effectiveness f_0. The minimum guarantees that the Kondo liquid susceptibility derived from the scaling formula does not exceed the experimental data during the numerical fit, which is always true as long as the Kondo liquid scaling holds. Once again, R_0 and r_l can be determined from the high temperature data with Eq. (4). We then fit the Hall coefficient in the intermediate temperature regime with two free parameters r_h and χ_0 by using

$$R_H = R_0 + r_l\rho(\chi - \chi_h) + r_h\chi_h. \qquad (6)$$

As an example, Fig. 4 shows the fit to the Hall data in Ce$_2$CoIn$_8$ [11]. The inset compares R_H with $\rho\chi$ calculated from experiment. The high temperature behavior of the Hall coefficient is well described by incoherent skew scattering. Below $T^* \approx 50\,$K, the prediction of Fert and Levy in Eq. (4) (dotted line) deviates severely from experiment. On the other hand, using the new formula and taking into account the coherent contribution, we obtain a good fit all the way down to $T_0 \approx 10\,$K, below which a ρ^2-dependence was claimed experimentally [11]. For comparison, Fig. 4 also plots the two contributions separately and the coherent skew scattering is seen to have a major contribution (dash-dotted line) in the Hall effect in Ce$_2$CoIn$_8$. In Fig. 1(b), the incoherent skew scattering is even more suppressed in the isostructural compound Ce$_2$PdIn$_8$ [15], giving rise to a similar χ_h-scaling discussed previously in Ce-115 compounds. In contrast, anomalous Hall effect in YbRh$_2$Si$_2$ [27] is dominated by incoherent skew scattering. While its coherence temperature is $\sim 70\,$K [28, 29], R_H follows the prediction of Eq. (4) for all temperatures above $\sim 7\,$K and only a slight change of slope (r_l) is seen at 90 K [27], possibly due to crystal field effect [30].

All together, while Ce$_2$CoIn$_8$ is a typical example whose anomalous Hall effect exhibits the two-component physics, Ce-115 and YbRh$_2$Si$_2$ represent the two rare extremes in realistic materials, in which either incoherent or coherent contribution is almost completely suppressed. Our results are summarized on a general phase diagram in Fig. 5. Above the characteristic temperature T^* determined by the onset of coherence [28], there are only localized f-moments and their incoherent skew scattering gives $R_H = R_0 + r_l\rho\chi$ predicted by Fert and Levy [6]. For $T < T_L$, all f-electrons become itinerant and the Hall effect is dominated by coherent skew scattering, for which the theory of Kontani and Yamada predicts



FIG. 4: (color online) Analysis of the Hall coefficient of Ce$_2$CoIn$_8$ [11]. The parameters are $T^* = 50\,$K, $R_0 = -4.12 \times 10^{-10}\,$m^3/C, and $\chi_0 = 5\,$memu/mol-Ce. For comparison, the dotted line shows the prediction of Fert and Levy with a constant r_l. The inset plots R_H versus $\rho\chi$.



FIG. 5: (color online) Anomalous Hall coefficient in different regimes on the temperature-pressure phase diagram.

$R_H \sim \rho^2$ [9]. Whether or not this ρ^2-scaling exists at extremely low temperatures depends on the details of the hybridization. In between, the f-electron states are described by the two-fluid physics so that both incoherent and coherent skew scatterings should be included, giving rise to the unusual two-component formula of Eq. (3). In the special case where incoherent skew scattering is completely suppressed, one finds a universal χ_h-scaling [16]. In the theory of Fert and Levy, the incoherent contribution originates from the interference between the f and d partial waves, with $r_l \propto \sin\delta_2 \cos\delta_2$, where δ_2 is the phase shift of the d-scattering channel [6]. Ce-115 [13, 14] and Ce_2PdIn_8 [15] fall into this category but the origin of the suppression has not been explained. A detailed study of the distinction between Ce_2CoIn_8 and Ce_2PdIn_8 may help resolve this issue.

In general, the specific conditions that control the relative importance of R_s^l and R_s^h in the two-fluid regime are not known. Although we were partly motivated by the theory of Kontani and Yamada [9], its validity for heavy electron materials remains unclear. A notable prediction of the theory is that the sign of r_h depends on the location of the resonant state around the Fermi energy so that Ce-compounds should have a positive r_h and Yb-compounds a negative r_h. Instead, we find a negative R_s^h and R_0 in Ce-115, Ce_2CoIn_8 and Ce_2PdIn_8 and a positive R_s^h and R_0 in $YbRh_2Si_2$, in line with the electron or hole nature of their charge carriers. A thorough understanding of the new formula may hence require a microscopic theory of the two-fluid physics that is not yet available. We will leave this for future work.

In conclusion, we propose an empirical formula for the anomalous Hall effect in heavy electron materials based on the phenomenological two-fluid framework. The new formula allows for a first consistent interpretation of the Hall experiment over a broad temperature range down to only a few Kelvin. It unifies the various scalings observed in different temperature regimes in experiment and provides a new basis for developing an improved theory incorporating previous theoretical proposals and the two-fluid physics. The identification of the incoherent and coherent contributions opens a new avenue for comparison between numerical calculations and experiment in the future. Investigations of similar phenomenon in other physical properties such as the spin Hall effect are yet another possibility.

The author thanks the stimulating discussion with D. Pines. This work is supported by NSF-China (Grant No. 11174339) and Chinese Academy of Sciences.

[1] N. Nagaosa *et al.*, Rev. Mod. Phys. **82**, 1539 (2010).
[2] R. Karplus and J. M. Luttinger, Phys. Rev. **95**, 1154 (1954).
[3] J. Smit, Physica **21**, 877 (1955).
[4] L. Berger, Physica **30**, 1141 (1964).
[5] P. Coleman, P. W. Anderson and T. V. Ramakrishnan, Phys. Rev. Lett. **55**, 414 (1985).
[6] A. Fert and P. M. Levy, Phys. Rev. B **36**, 1907 (1987).
[7] H. Kohno and K. Yamada, J. Magn. Magn. Matter. **90&91**, 431 (1990).
[8] K. Yamada, H. Kontani, H. Kohno and S. Inagaki, Prog. Theor. Phys. **89**, 1155 (1993).
[9] H. Kontani and K. Yamada, J. Phys. Soc. Jpn. **63**, 2627 (1994).
[10] U. Köhler *et al.*, J. Phys.: Condens. Matter **19**, 386207 (2007).
[11] G. Chen *et al.*, J. Phys.: Condens. Matter **15**, S2175 (2003).
[12] Y.-F. Yang *et al.*, J. Phys.: Conf. Ser. **273**, 012066 (2011).
[13] M. F. Hundley, A. Malinowski, P. G. Pagliuso, J. L. Sarrao and J. D. Thompson, Phys. Rev. B **70**, 035113 (2004).
[14] Y. Nakajima *et al.*, J. Phys. Soc. Jpn. **76**, 024703 (2007).
[15] D. Gnida, M. Matusiak and D. Kaczorowski, Phys. Rev. B **85**, 060508(R) (2012).
[16] Y.-F. Yang and D. Pines, Phys. Rev. Lett. **100**, 096404 (2008).
[17] Y.-F. Yang and D. Pines, arXiv:1206.1115, unpublished.
[18] J. Schoenes, C. Schönenberger, J. J. M. Franse and A. A. Menovsky, Phys. Rev. B **35**, 5375(R) (1987).
[19] J. H. Shim, K. Haule and G. Kotliar, Science **318**, 1615 (2007).
[20] Y.-F. Yang, Phys. Rev. B **79**, 241107(R) (2009).
[21] V. Barzykin and D. Pines, Adv. Phys. **58**, 1 (2009).
[22] O. O. Bernal *et al.*, Physica B **281&282**, 236 (2000).
[23] K. R. Shirer *et al.*, arXiv:1206.1879, unpublished.
[24] S. Nakatsuji, D. Pines, and Z. Fisk, Phys. Rev. Lett. **92**, 016401 (2004).
[25] N. J. Curro, B.-L. Young, J. Schmalian and D. Pines, Phys. Rev. B **70**, 235117 (2004).
[26] J. A. Mydosh and P. M. Oppeneer, Rev. Mod. Phys. **83**, 1301 (2011).
[27] S. Paschen *et al.*, Physica B **44-46**, 359 (2005).
[28] Y.-F. Yang *et al.*, Nature **454**, 611 (2008).
[29] S.-K. Mo *et al.*, Phys. Rev. B **85**, 241103(R) (2012).
[30] S. Ernst *et al.*, Nature **474**, 362 (2011).